

872 W. Baltimore Pike, PO Box 9, West Grove, PA 19390 800-367-5296, 610-869-4024, FAX: 610-869-0171 cuserv@jacksonimmuno.com, www.jacksonimmuno.com www.jireurope.com

PRODUCT SPECIFICATIONS

Product: Peroxidase-conjugated AffiniPure Goat Anti-Rat IgM, µ Chain Specific

Code Number: 112-035-020

Lot Number: 68900

Physical State: Freeze-dried powder

Size: 2.0 ml

Antibody Concentration: 0.8 mg/ml

Suggested Dilution Range: 1:500 - 1:5,000 for immunohistochemistry on tissue sections

1:5,000 - 1:100,000 for ELISA and Western blotting with chromogenic substrates

1:10,000 - 1:200,000 for Western blotting with ECL substrates

Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6

<u>Stabilizer</u>: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)

<u>Preservative</u>: None (Warning: Use of sodium azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase.)

Note: Due to tightened regulations in many countries and states, thimerosal, a mercury-containing preservative, will no longer be included in our products. Please follow our recommendation for storage to avoid microbial contamination.

Reconstitution and Storage: Store freeze-dried product at 2-8°C until reconstitution. When ready to use, reconstitute with 2.0 ml d. water. Centrifuge product if the solution is not clear after reconstitution. To judge clarity, draw product into a clean pasteur pipette. Storing the product for more than one day at final working dilutions is not recommended. For extended storage as a liquid, add an equal volume of glycerol (ACS or better grade) for a final concentration of 50% followed by storage at -20°C. Please note that the concentration of protein and buffer salts will be one-half of the original after the addition of glycerol. Alternatively, the product may be aliquoted and frozen at -70°C or below, in the absence of glycerol. It is important to avoid repeated freezing and thawing. Expiration date: one year from date of reconstitution.

<u>Purity</u>: The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.

<u>Antibody Specificity</u>: Based on immunoelectrophoresis, the antibody reacts with the heavy chain portion of rat IgM. No antibody was detected against IgG or non-immunoglobulin serum proteins, but the antibody may cross-react with IgM from other species.

Country of Origin: USA

Note: For in vitro research use only, not for diagnostic or therapeutic use. This product is not a medical device.